

LA-UR-21-29053

Approved for public release; distribution is unlimited.

Title: Coupled radiation transport and plasma physics for ICF simulations

Author(s): Hammer, Hans Ruediger
Park, HyeongKae
Chacon, Luis
Taitano, William
Anderson, Steven Edward
Simakov, Andrei N.

Intended for: Presentation to external collaborators

Issued: 2021-09-14

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Coupled radiation transport and plasma physics for ICF simulations


Hans Hammer HyeongKae Park Luis Chacón William Taitano
Steven Anderson Andrei Simakov

March 9, 2021

Motivation

Describing high-temperature plasmas requires inclusion of several nonlinear coupled physics

- Ion kinetics - for each isotope species present in the plasma
- Electron kinetics
- Thermal radiative transfer (TRT)
- Electromagnetic description
- Nuclear reactions (Fusion burn)



TRT is a key physics component in ICF

- Thermal x-ray radiation becomes the dominant energy transfer mechanism in hohlraum
- Driver of capsule implosion
- Radiative energy loss

Scope of this project

Simulating plasmas in ICF hohlraums.

Our project aims to solve

- Kinetic ions (particles)
- Fluid electrons
- Energetic electrons
- Electromagnetics
- Kinetic radiation (MG DP solver)
- Laser-plasma interactions

Coupling of the different physics is done in a nonlinear LO system

- Moment (fluid) equations for plasma
- Gray radiation diffusion
- Consistency terms enforce HO solution
- Concept has been tested in iFP

This presentation describes work done in iFP, which is slightly different



DR project equations - Overview

We try to solve a coupled system with these equations
(missing hot electrons, laser-plasma-interactions):

Ion Vlasov-Fokker-Planck:
$$\partial_t f_\alpha + \mathbf{u} \cdot \nabla_x f_\alpha + \frac{q_\alpha}{m_\alpha} \mathbf{E} \cdot \nabla_u f_\alpha = \sum_\beta C_{\alpha\beta}$$

Radiation transport:
$$\frac{1}{c} \partial_t I + \Omega_i \partial_i I + \sigma_t I - \sigma_e B = S_{\Omega' \rightarrow \Omega, \nu' \rightarrow \nu}$$

Quasineutrality:
$$\sum_\alpha q_\alpha n_\alpha + q_e n_e = 0$$

Ambipolarity:
$$\sum_\alpha q_\alpha n_\alpha u_{\alpha,i} + q_e n_e u_{e,i} = 0$$

Electron temperature
$$\partial_t \frac{3}{2} n_e k_B T_e + \partial_i u_{e,i} \left[\frac{3}{2} n_e k_B T_e + P_e \right] + \partial_i Q_{e,i} - q_e n_e u_{e,i} E_i - \sum_\alpha W_{e\alpha} - S_{re} = 0$$

Ohm's law:
$$E_i = \frac{\partial_i P_e + \sum_\alpha F_{\alpha e,i} - S_{rp,i}}{q_e n_e}$$



Vlasov-Fokker-Planck equation

Electrostatic kinetic plasma equation with Coulomb collisions

$$\partial_t f_\alpha + \mathbf{u} \cdot \nabla_x f_\alpha + \frac{q_\alpha}{m_\alpha} \mathbf{E} \cdot \nabla_u f_\alpha = \sum_\beta C_{\alpha\beta}$$

where

- $f_\alpha(\mathbf{x}, \mathbf{u}, t)$ velocity distribution function for species α
- $\mathbf{E}(\mathbf{x})$ electric field
- q_α species charge
- m_α species mass
- $C_{\alpha\beta}(f_\alpha, f_\beta)$ Fokker-Planck collision operator with species β (Rosenbluth et al., 1957)
- ∇_x gradient operator in physical space
- ∇_u gradient operator in velocity space



Thermal Radiative Transfer Equation

Thermal radiative transfer (TRT) with physical scattering can be described by the following equation

$$\frac{1}{c} \partial_t I + \boldsymbol{\Omega} \cdot \nabla_{\mathbf{x}} I + \sigma_t I - \sigma_e B = \int_0^\infty \int_{4\pi} \sigma_s(\boldsymbol{\Omega}' \rightarrow \boldsymbol{\Omega}, \nu' \rightarrow \nu) I(\boldsymbol{\Omega}', \nu') d\boldsymbol{\Omega}' d\nu'$$

where

- $I(\mathbf{x}, \boldsymbol{\Omega}, \nu, t)$ Radiation intensity in direction $\boldsymbol{\Omega}$ and frequency ν
- $B(\nu, T_e)$ Planck function (nonlinear emission spectrum)
- $\sigma(\mathbf{x}, \rho, \nu, T_e)$ Opacity (total, emission, and scattering)

Nonlinearly coupled to electron physics by ρ and T_e . The frequency-integrated emission source is $\propto T_e^4$




Common Approach for Radiation-Plasma coupling

This system is difficult to solve

- Stiff coupling between radiation and material temperature ($\propto T^4$)
- High dimensionality of the transport equation

Common approaches to solve this

- Linearization of emission-source
- Operator-split temporal update
- Monte-Carlo methods for Boltzmann integral



ICF Hohlraum. Source: LLNL

State of the Art in Plasma-Radiation coupling

Current radiation-hydrodynamics codes use a fluid model for plasma

- Plasma described by bulk quantities (moments)
- Assume the plasma is sufficiently collisional (Small Knudsen number)
- Heat flux limiter required to avoid nonphysical behavior


These assumptions are violated in ICF plasmas

- Free streaming of ions and electrons
- Counter-flows
- Non-maxwellian velocity distribution

We need to understand when and where these assumptions are not valid, and the effects



Multi-physics coupling through low-order system



To efficiently couple these equations, we employ a system of low-order (LO) equations

- LO system is tightly coupled
 - Moments of the kinetic equations
 - Implicit time-discretization
 - Solved with non-linear solver (e.g. Newton-Krylov)
- LO-equations are corrected by high-order (HO) solution
 - Example of existing methods: Variable Eddington tensor formulation/ quasi-diffusion
- HO-solvers are (mostly) independent from each other
 - HO-solvers use LO-solution for coupled parameters (e.g., T_e for emission source, u_e for motion correction)

Kinetic electrons have been implemented in a different branch of the iFP code



LO equations - Ions (Moments of VFP)

Conservation of mass (continuity)

$$\partial_t \rho_\alpha + \partial_i \rho_\alpha u_{\alpha,i} + \gamma_{V,\rho,\alpha} - \gamma_{FP,\rho,\alpha} = 0,$$

Conservation of momentum

$$\partial_t p_{\alpha,i} + \partial_j p_{\alpha,i} u_{\alpha,j} + \partial_i P_\alpha - q_\alpha n_\alpha E_i + \sum_\beta F_{\alpha\beta,i} + F_{\alpha e,i} + \gamma_{V,p,\alpha,i} - \gamma_{FP,p,\alpha,i} = 0,$$

Conservation of energy

$$\partial_t U_\alpha + \partial_i u_{\alpha,i} [U_\alpha + P_\alpha] - q_\alpha n_\alpha u_{\alpha,i} E_i + \sum_\beta W_{\alpha\beta} + W_{e\alpha} + \gamma_{V,U,\alpha} - \gamma_{FP,U,\alpha} = 0.$$

where

- Mass density $\rho_\alpha = m_\alpha n_\alpha$
- Bulk velocity $u_{\alpha,i}$ (Tensor notation)
- Total energy $U_\alpha = 3/2 n_\alpha k_B T_\alpha + 1/2 \rho_\alpha u_{\alpha,i}^2$
- Friction terms $F_{\alpha\beta}$
- Energy exchange terms $W_{\alpha\beta}$
- Pressure P_α
- Consistency terms γ_V and γ_{FP}



LO equations - Electrons (Hybrid)

The electron number density can be calculated using quasi-neutrality

$$\sum_{\alpha} q_{\alpha} n_{\alpha} + q_e n_e = 0,$$

and the electron velocity using ambipolarity

$$\sum_{\alpha} q_{\alpha} n_{\alpha} u_{\alpha,i} + q_e n_e u_{e,i} = 0.$$

The electron temperature equation is coupled with radiation

$$\partial_t \frac{3}{2} n_e k_B T_e + \partial_i u_{e,i} \left[\frac{3}{2} n_e k_B T_e + P_e \right] + \partial_i Q_{e,i} - q_e n_e u_{e,i} E_i - \sum_{\alpha} W_{e\alpha} - S_{re} = 0,$$

where

- Heat flux $Q_{e,i}$
- Radiation energy deposition S_{re}



LO equations - Electric field (Ohm's Law)

Using the electron momentum equation

$$\partial_t p_{e,i} + \partial_j [u_{e,j} p_{e,i}] + \partial_i P_e - q_e n_e E_i + \sum_{\alpha} F_{\alpha e,i} - S_{rp,i} = 0,$$

and assuming $m_e \approx 0 \rightarrow p_{e,i} \approx 0$ to get Ohm's law:

$$E_i = \frac{\partial_i P_e + \sum_{\alpha} F_{\alpha e,i} - S_{rp,i}}{q_e n_e},$$

with $S_{rp,i}$ the radiation momentum deposition.

The DR will use a generalized version of this electric field including electromagnetic effects, Andrei is working on a rigorous formulation



LO equations - Radiation (Gray Diffusion)

Radiation energy conservation

$$\partial_t E_r + \partial_i F_i + S_{re} = 0,$$

Radiation momentum conservation

$$\frac{1}{c^2} \partial_t F_i + \partial_j P_{r,ij} + S_{rp,i} = 0,$$

where

- Gray radiation energy density $E_r = \frac{1}{c} \int_0^\infty \int_{4\pi} I d\Omega d\nu$
- Gray radiative flux $F_i = \int_0^\infty \int_{4\pi} \Omega_i I d\Omega d\nu$
- Radiation pressure $P_{r,ij} = \int_0^\infty \int_{4\pi} \Omega_i \Omega_j I d\Omega d\nu$



Radiation - Plasma coupling

Radiation and plasma couple through energy and momentum exchange terms.

Energy

$$S_{re} = \sigma_a c E_0 - \sigma_e a c T_e^4 + \frac{\sigma_t}{c} u_{e,i} F_{0,i},$$

Momentum (second term negligible for non-relativistic regimes)

$$S_{rp,i} \simeq \frac{\sigma_t}{c} F_{0,i} + \frac{u_{e,i}}{c^2} (\sigma_a c E_0 - \sigma_e a c T_e^4) \approx 0$$

Isotropic scattering is hidden in the opacities: $\sigma_t = \sigma_s + \sigma_a$

Radiation is assumed to only interact with electrons. Coupling is dependent on u_e and T_e



Radiation - Material motion correction

Interaction of radiation with moving material require material motion corrections

- Using comoving frame
- Lorentz transformation of radiation quantities
- Emission and scattering (Thompson) are isotropic in comoving frame
- Becomes anisotropic in laboratory frame

Simplified material motion correction for non-relativistic velocities (Morel, 2006):

$$F_{0,i} = F_i - u_{e,i}E_r - u_{e,j}P_{r,ij}$$

$$E_0 \simeq E_r - 2 \frac{u_{e,i}}{c^2} F_{0,i} \approx 0$$

$$P_{r,ij} = \mathcal{E}_{ij}E_r \quad \text{with} \quad \mathcal{E}_{ij} \simeq \frac{1}{3}\delta_{ij} \quad (\text{Eddington tensor})$$



Radiation Diffusion Approximation

Neglecting the temporal term in the radiation momentum equation, and with the non-relativistic material motion corrections, we get

$$F_i = -\frac{c}{\sigma_t} \partial_j \mathcal{E}_{r,ij} E_r + u_{e,i} E_r + u_{e,j} \mathcal{E}_{ij} E_r,$$

and a simplified momentum deposition

$$S_{rp} = -\partial_j [\mathcal{E}_{ij} E_r].$$

Cancellations

Several terms are required to cancel

- Stem from different terms and show up in different equations
- Discretization must match



Radiation-Hydrodynamic asymptotic limit

The previous equations yield the rad-hydro formulation in the highly collisional limit

- Substitute radiation coupling terms and cancel
- The plasma fluid model reduces to Euler equations

Radiation diffusion equation simplifies to

$$\partial_t E_r - \partial_i \frac{c}{\sigma_t} \partial_i [\mathcal{E} E_r] + \partial_i [u_{e,i} (1 + \mathcal{E}) E_r] + \sigma_a c E_0 - \sigma_e a c T_e^4 - u_{e,i} \partial_i [\mathcal{E} E_r] = 0,$$

which gives with the chain-rule the CRASH equation (Holst et al., 2011)

$$\partial_t E_r - \partial_i \frac{c}{\sigma_t} \partial_i [\mathcal{E} E_r] + \partial_i [u_{e,i} E_r] + \mathcal{E} E_r \partial_i u_{e,i} + \sigma_a c E_0 - \sigma_e a c T_e^4 = 0$$



Verification M45 Radiative Shock

Planar Mach 45 shock problem (Lowrie and Edwards, 2008)

- Shock driven by radiation
- Self-similar solution
- Strongly collisional regime (iFP should reproduce rad-hydro)


Input is for rad-hydro, must be translated to plasma parameters. Special care is required

- It looks like a proton (Hydrogen) plasma ($m \approx 1$)
- Rad-hydro input neglected electrons for heat capacity (error)
- To match both heat capacity (sum ions and electrons) and mass density, ion mass must be doubled ($m \approx 2$) (fix)


Special care must be taken for rad-hydro inputs.



Radiative Shock - Results




(a) Mass density




(b) Velocity



Radiative Shock - Results II



(c) Temperature



(d) Temperature at shock peak

Conclusions

Implemented radiation-plasma coupling in the iFP LO solver

- Currently Gray radiation diffusion
- Verified with radiative Mach 45 shock
- Holds in the asymptotic limit

This work is the foundation to ground-breaking improvements

- Extension with HO solver (multi-group diffusion, transport)
- More sophisticated opacity models (tables, LTE, NLTE)

Radiation coupling lessons can be directly translated to DR project



References



Holst, B. van der et al. (May 2011). “CRASH: A BLOCK-ADAPTIVE-MESH CODE FOR RADIATIVE SHOCK HYDRODYNAMICS—IMPLEMENTATION AND VERIFICATION”. In: *The Astrophysical Journal Supplement Series* 194.2, p. 23. ISSN: 0067-0049. DOI: 10.1088/0067-0049/194/2/23.



Lowrie, Robert B. and Jarrod D. Edwards (July 1, 2008). “Radiative Shock Solutions with Grey Nonequilibrium Diffusion”. In: *Shock Waves* 18.2, pp. 129–143. ISSN: 1432-2153. DOI: 10.1007/s00193-008-0143-0.



Morel, Jim E. (2006). “Discrete-Ordinates Methods for Radiative Transfer in the Non-Relativistic Stellar Regime”. In: *Computational Methods in Transport*. Ed. by Frank Graziani. Lecture Notes in Computational Science and Engineering. Springer Berlin Heidelberg, pp. 69–81. ISBN: 978-3-540-28125-2.



Rosenbluth, Marshall N., William M. MacDonald, and David L. Judd (July 1, 1957). “Fokker-Planck Equation for an Inverse-Square Force”. In: *Physical Review* 107.1, pp. 1–6. DOI: 10.1103/PhysRev.107.1.

